arXiv:astro-ph/0202065v1 4 Feb 2002

TWO YEARS OF THE X-RAY SKY IN M31

A.K.H. Kong, M.R. Garcia, F.A. Primini, R. Di Stefano, and S.S. Murray

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, U.S.A.

Abstract

We have been monitoring M31 with HRC and ACIS onboard *Chandra* regularly in the past two years. By combining eight *Chandra* ACIS-I observations taken between 1999 to 2001, we have identified 204 X-ray sources within the central $\sim 17' \times 17'$ region of M31, with a detection limit of $\sim 1.6 \times 10^{35}\,\mathrm{erg\,s^{-1}}$. Of these 204 sources, 21 are identified with globular clusters, 2 with supernova remnants (one of them is spatially resolved with *Chandra*), and 8 with planetary nebula. By comparing individual images, about 50% of the sources are variable in time scales of months. We also found 14 transients. Combining all the available transients found in literatures, there are 25 transients in M31 and M32 detected by *Chandra* and *XMM-Newton*; we present some of the long-term lightcurves by using the HRC, ACIS and *XMM-Newton* data. The spectral shape of 12 sources is shown to be variable, suggesting that they went through state changes. The luminosity function of all the point sources is consistent with previous observations (a broken power-law with a luminosity break at $1.7 \times 10^{37}\,\mathrm{erg\,s^{-1}}$). However, when the X-ray sources in different regions are considered separately, different luminosity functions are obtained. This indicates that the star formation history might be different in different regions.

Key words: galaxies: individual (M31) – X-rays: galaxies

1. Introduction

M31 was observed by *Chandra* and *XMM-Newton* soon after these observatories were launched. In the first observation (8.8 ks) of the core region by *Chandra* in 1999 October, 121 point sources were identified within the central $17' \times 17'$ region and the nucleus was nicely resolved into five point sources which were not seen in previous missions (Garcia et al. 2000a). Moreover, a bright transient was discovered $\sim 26''$ from the nucleus. A relatively deeper *XMM-Newton* observation (34.8 ks) was made in 2000 June; 116 sources were detected down to a limiting luminosity of $6 \times 10^{35}\ \mathrm{erg\,s^{-1}}$ (0.3–12 kpc; Shirey et al. 2001) and a pulsating supersoft transient with a periodicity of ~ 865 s was discovered (Osbrone et al. 2001). Moreover, both *Chandra* (Garcia et al. 2001a; Primini et al. 2000) and *XMM-Newton* (Shirey et al. 2001) observations confirmed that the unresolved X-ray emission in the core region is much softer than most of the resolved X-ray sources in that region. Fifteen X-ray point sources are newly associated with globular clusters (Di Stefano et al. 2002), which when combined with the previous *ROSAT* results (Supper et al. 2001) brings the total number of M31 globular clusters with detected X-ray emission to 48. *Chandra* and *XMM-Newton* also discovered several bright ($L_X > 10^{37}$ $\mathrm{erg\,s^{-1}}$) transients in M31 and M32 (Garcia et al. 2000a; Garcia et al. 2000b; Osbrone et al. 2001; Shirey et al. 2001; Kong et al. 2001; Garcia et al. 2001b). The brightest of these reached a peak luminosity of $L_X \sim 3 \times 10^{38}$ $\mathrm{erg\,s^{-1}}$(Kong et al. 2001).

We report herein a brief summary of our *Chandra* monitoring program of M31 in the past two years; it will mainly focus on the properties (identifications, temporal and spectral variability, and luminosity function) of point sources in the central $\sim 17' \times 17'$ region of M31 as deduced from eight separate ~ 5ks ACIS-I observations spanning ~ 1.5 years. In addition, we will also present early results of an X-ray resolved supernova remnant and an updated “X-ray movie” of M31 up to 2002 January.

2. Observations

M31 was observed with *Chandra* regularly as part of the AO-1 and AO-2 GTO program during 1999–2001 (This program continues as GO observations in AO-3). The program is originally designed to search for transients in M31. These observations consist of a series of HRC snapshots ($\sim$ 1 ks) which cover the whole galaxy (see Figure 1); once a transient is found, a follow-up ACIS ($\sim$ 5 ks) observation then images the newly-discovered transient, otherwise, ACIS points to the nuclear region. In this paper, we mainly focus on the ACIS-I (I0, I1, I2 and I3) data centered in the central $16.9' \times 16.9'$ region of M31; this consists of 8 separate observations from 1999 October to 2001 June, with exposure times ranging from 4 to 8.8 ks. The actual region covered by the *Chandra* observations is slightly larger than $16.9' \times 16.9'$ due to different roll angles. Sources near the outer edge of ACIS are therefore not observable in all eight exposures. In addition, we also make use of the HRC data and other ACIS pointings to construct the long-term lightcurves of sources.

In order to create a deep image suitable for the detection of faint sources, the eight observations were combined

Proc. Symposium *‘New Visions of the X-ray Universe in the XMM-Newton and Chandra Era’* 26–30 November 2001,



Figure 1. The field-of-view of HRC-I (blue) and ACIS-I (red) pointings overlaid on an optical Digitized Sky Survey image of M31. The aim-points are marked with a cross.

into a single stacked image with a total integration time of 39.7 ks. Figure 2 shows the stacked "true color" image of M31, which is a composite of images from soft (0.3–1.0 keV), middle (1–2 keV) and hard (2–7 keV) bands. Soft sources appear red, moderately hard sources appear green, and the hardest sources appear blue. The image has been corrected for exposure and smoothed slightly with a Gaussian ($\sigma = 0.5''$) in order to improve the appearance of point sources; diffuse emission around the nucleus is also clearly shown. Also shown in Figure 2 is the "true color" image of the central $2' \times 2'$ region of M31 with $1/8''$ pixel resolution, with the possible nuclear counterpart (M31*) marked (Garcia 2001).

3. Source Detection and Identification

Discrete sources in the stacked image were found with WAVDETECT (Freeman et al. 2002), a wavelet detection algorithm implemented within CIAO. The central $2' \times 2'$ region was treated separately by using the $1/8''$ image. A total of 204 sources were detected above 2.5σ.

We cross-correlated the source list with existing catalogs of M31 objects. We find that 77 *Chandra* sources have counterparts in the *ROSAT* HRI catalog (Primini et al. 1993). The remaining 127 (= 204 − 77) *Chandra* sources were not detected in the *ROSAT* HRI catalog, presumably because they are below the *ROSAT* detection limit or are variable. This *Chandra* catalog extends $\sim 5\times$ fainter than the *ROSAT* HRI catalog. We identify 22 *Chandra* sources with globular clusters; twelve of these globular clusters are identified as X-ray sources for the first time. We find two matches with supernova remnant (SNR). In particular, one of the them is spatially resolved by *Chandra* (see § 5). Eight planetary nebula (PN) are found to be associated with our *Chandra* sources. However, the X-ray luminosity of our PN candidates are at least 3 order of magnitude brighter than that in our Galaxy (e.g. Kastner et al. 2001); the X-ray colors are generally harder as well. Therefore, it is possible that these PN are either very unusual or something other than PN. We suggest that these may be objects similar to GX 1+4 in our Galaxy (i.e. symbiotic stars with a neutron star companion). We searched for matches between stellar nova as listed in the IAUC contemporaneous to and within the field-of-view of our *Chandra* observations, and found no matches within $3''$. We also searched for matches with OB associations, as O and B stars may be moderate X-ray sources ($L_X < 10^{33}$erg s^{-1}, Berghoefer et al. 1997). While this is well below our detection limit, a group of O and/or B stars may reach our detection threshold, and star forming regions could conceivably harbor massive X-ray binaries. However, we found no matches within our search radius of $3''$. Five possible foreground objects are also found in our field based on their X-ray colors and coincidence of stars.

4. Temporal and Spectral Variability

The eight *Chandra* ACIS-I observations described herein span nearly 2 years from 1999–2001. This is substantially longer than previous surveys by *ROSAT* (2 observations separated by ~ 1 year). In order to study long-term X-ray variability, we computed a variability parameter following Primini et al. (1993). We found 99 X-ray variables, corresponding to $\sim 50\%$ of the total. By comparing *ROSAT* observations to *Einstein* observaitions 10 years earlier, Primini et al. (1993) found that $\sim 42\%$ of the X-ray sources within central $7.5'$ region were variable. By comparing two *XMM-Newton* observaitions separated by six months Osbrone et al. (2001) found that $> 15\%$ of the sources in the central $30'$ were variable.

We have also discovered 13 bright transients for which the source is found in at least one observation with luminosity of $\gtrsim 5 \times 10^{36}$ erg s^{-1} and is not detected in at least one of the other observations. Note that the luminosity limit covers typical outburst luminosities of soft X-ray transients and Be/X-ray binaries in our Galaxy. One important transient was missed by this analysis because it had a peak luminosity below our "bright transient" threshold during the eight ACIS-I observations considered here. This object is XMMU J004234.1+411808 (= CXOM31



Figure 2. Top: Stacked "true color" Chandra ACIS-I image (39.7 ks) of the central $\sim 17' \times 17'$ *region of M31. This image was constructed from the soft (red; 0.3–1 keV), medium (green; 1–2 keV) and hard (blue; 2–7 keV) energy bands. The pixel size is* $1.96''$ *and the image has been smoothed with a Gaussian (*$\sigma = 0.5''$*) function. Bottom: Stacked "true color" Chandra ACIS-I image of the central* $\sim 2' \times 2'$ *region of M31. The pixel size is* $0.123''$*. M31* candidate is marked.*



Figure 3. Lightcurves of 5 bright transients and globular cluster Bo 86 as seen in the past two years with Chandra and XMM-Newton. Observations from HRC-I (blue), ACIS-I (red), ACIS-S (green) and XMM-MOS/PN (yellow) are included.



Figure 4. Spectral changes of three of the sources. The first two from the left change from lower luminosity at softer state to higher luminosity with harder state. The bright transient in the right shows a harder spectrum during its decay from the peak of the outburst.

J004234.3+411809), which Osbrone et al. (2001) suggest is an X-ray nova. By adding all the published X-ray transients in M31 found by *Chandra* and *XMM-Newton* (Osbrone et al. 2001; Shirey 2001; Kong et al. 2001; Garcia et al. 2001b; Trudolyubov et al. 2002), we conclude that there are 25 "bright" transients in M31 (and M32) in the past two years.

Figure 3 shows five representative lightcurves of transients found in M31 and M32. The transient discovered with *Chandra* in the first observation (Garcia et al. 2000a) remained in outburst for more than one year and finally turned off in 2001 June (Figure 3). In addition, three HRC-I data points indicate that the source underwent flarings or state transitions during the early stage of the outburst. During the whole outburst, the energy spectrum is consistent with a power-law with slope of $\sim$ 1.5 (see inset of Figure 3). The remaining three transients show typical lightcurves with a fast rise followed by an exponential decay like those seen in our Galaxy (Chen et al. 1997). The characteristic decay time is about 30–50 days. CXOM31 J004309.7+411901 is a recurrent transient; the source was detected with *Einstein*, *ROSAT* HRI and *ROSAT* PSPC at luminosities ranging from 1 to 8 $\times 10^{37}$ erg s^{-1}. Supper et al. (2001) noted variability between PSPC exposures of a factor of 5. Except for the first *Chandra* observations, the source underwent a state transition to the low state at $L_X < 10^{36}$ erg s^{-1}; it re-brightened to 3×10^{37} erg s^{-1}in 2001 November and shows decay in 2002 January observations. Beginning in the summer of 2001, we also monitored newly discovered X-ray transients with *HST* to search for UV counterparts. This program allows us to follow the evolution of six transients in M31 during their outbursts in both X-rays and UV; it will be the first step in extending the highly successful *RXTE*/ASM and optical follow-up of Galactic X-ray transients to our nearest neighbor spiral galaxy.

In addition to transients, we also found several sources which have interesting long-term lightcurve. For example, CXOM31 J004218.5+411758 is in the M31 globular cluster Bo 86 and shows a possible $\sim$ 200 day modulation. This is reminiscent of the Galactic source 4U 1820–30 in the globular cluster NGC 6624, which has a 176-d long-term modulation. It is worth noting that X-ray variability on long time-scales (from days to years) has been found in many low-mass and high-mass X-ray binaries in our Galaxy (see e.g. Kong & Charles 2000).

We searched for spectral variability in all of the sources using a method analogous to that described in Primini et al. (1993), but replacing the flux with hardness ratio. Only 12 sources show spectral variability, corresponding to 6% of the total population. This is of course a lower limit, as small changes in the spectra of weak sources are undetectable with the number of counts accumulated in our 40 ks of merged data. In order to further investigate the nature of the spectral variations, we fit simple spectra to two of the brighter spectral variables. The fits show that as the counting rate increases, the spectrum becomes harder (see Figure 4). This is reminiscent of atoll and Z sources in our Galaxy (see e.g. Hasinger & van der Klis 1989). The luminosity of these two sources ranges from $(0.4 - 1.0) \times 10^{38}$ erg s^{-1}, which is higher than the typical luminosity ($< 10^{37}$ erg s^{-1}) of atoll sources. However, this luminosity is similar to that of the Z sources, which are believed to reach the Eddington limit (Psaltis et al. 1995). The luminosity and spectral changes appear consistent with that seen in Z sources as they move along the "normal branch". These are probably the first extragalactic Z sources to be identified, except for LMC X–2 (Smale & Kuulkers 2000). Continued monitoring may confirm the nature of these sources by allowing us to trace out the full Z-shape of the spectral variations expected if this analogy is correct. Spectral variation is also seen in transients dur-



Figure 5. Left: "True color" Chandra ACIS-I image of CXOM31 J004327.7+411829. This image was constructed from the soft (red: 0.5–1 keV), medium (green: 1–2 keV) and hard (blue: 2–7 keV) energy bands. The pixel size is $0.496''$ *and the image has been smoothed with a* $0.496''$ *FWHM Gaussian function. Right: Hardness-ratio (1–2 keV over 0.5–1 keV) image of the SNR, with X-ray contours overlaid. The contour levels are arbitrarily chosen (from 0.5–7 keV smoothed image) to best follow the surface brightness variations. Note the "hot spots" in the north-western arc and southern region of the SNR.*

ing their evolution away from the outburst. Figure 4 shows the spectral evolution of the brightest transient ($\sim 3\times10^{38}$ erg s^{-1}) ever found in M31 (Kong et al. 2001); the source becomes harder during the decay of the outburst. While the source is in its decay stage, it is still relatively bright in a recent observations taken in 2002 January ($L_X \gtrsim 10^{37}$ erg s^{-1}).

5. Discovery of an X-ray Resolved SNR

We have identified two SNRs in the central region of M31 with *Chandra*, and both of them were detected by *ROSAT* previously. One of them (CXOM31 J004327.7+411829) is found to be extended in X-rays as a distinct ring-shaped object with a diameter of $\sim 11''$ (33 pc). Figure 5 shows the "true color" X-ray image of the SNR; also shown is the hardness ratio image of the SNR. It is clear that bright spots can be seen in north-western and southern regions of the SNR. This SNR is also detected in several optical surveys (e.g. d'Odorico et al. 1980; Blair et al. 1981); it was identified as an irregular, faint and high [S II]/H$_\alpha$ (~ 1) SNR. The optical extension is about $10'' - 18''$, with a crescent open in the south-eastern part of the SNR. Our X-ray image also shows similar morphology.

Initial examination of a 40ks ACIS observations on 2001 October indicates that a non-equilibrium ionization (NEI) model with interstellar absorption is appropriate for modeling the SNR's spectrum. It is also evident that the spectrum is dominated by a broad emission of O VIII line at 0.654 keV and a blend of Fe L shell lines and Ne K shell lines around 0.9 keV. The best fitting NEI model indicates a 0.3–7 keV emitted luminosity of $\sim 5 \times 10^{36}$ erg s^{-1}.

6. Luminosity Function

In Figure 6 we plot the cumulative luminosity function (CLF) for all detected sources in the stacked image, and also separately the LFs of inner bulge (region 1; central $2' \times 2'$), outer bulge (region 2; central $8' \times 8'$ excluding region 1), disk (region 3; central $17' \times 17'$ excluding regions 1+2), and bulge (regions 1+2 combined). The LF for all sources has a break at $\sim 2 \times 10^{37}$ erg s^{-1}, with $\alpha_1 = 0.38$ before the break and $\alpha_2 = 1.63$ after the break. This result is in good agreement with previous *ROSAT* (Primini et al. 1993) and *XMM-Newton* (Shirey et al. 2001) measurements of the LF.

The CLF for the inner bulge is significantly different, with a break at a lower luminosity ($\sim 1.6 \times 10^{36}$) and a significantly flatter distribution at the faint end. We performed a two-sample Kolmogorov-Smirnov (K-S) test for the LFs of inner bulge and disk, and find that there is only 3% probability that they are drawn from the same distribution. Simulations show that the flattening of the CLF of inner bulge is not due to incompleteness.

However, the counts in a CLF are not independent and therefore direct fits to it will underestimate the errors and may produce a biased best estimate of the slope. In order to more accurately estimate the errors and slopes, we used a maximum likelihood method (e.g. Crawford et al.



Figure 6. Left: Luminosity functions for all sources (Regions 1+2+3) and bulge (Regions 1+2). Right: Luminosity functions for inner bulge (Region 1), outer bulge (Region 2) and disk (Region 3).

1970) to determine the slopes in the differential luminosity functions (DLFs). For the inner bulge, the LF is roughly consistent with a single power-law with $\alpha \sim 0.7$ although the appearance of the LFs (Figure 6) and the results of simulations indicate the presence of a brake to a flatter LF at the lowest fluxes. However, we note that the statistics in the inner bulge region are insufficient to confirm the presence and constrain the size of the break.The slope of the outer bulge and disk below the break is roughly consistent with 0.5–0.6. Further away from the inner bulge, a break appears near 10^{37} erg s^{-1} and it shifts to higher luminosity when considering the disk region (see Figure 6). In addition, the slope becomes steeper (from $\sim$ 1.0 to 2.2) when moving from the bulge to disk.

The black holes and neutron stars that power many of the M31 X-ray sources have formed through the evolution of initially massive stars. Because of this, the X-ray LF traces the history of star formation and evolution of these massive stars in binary systems. Breaks in the LF may indicate indicate an impulsive star formation event. As the X-ray binaries age, their average luminosity shifts to lower values and therefore the location of the break may be an indication of the how long ago the star formation event occurred (Wu et al. 2002). Luminosity functions which do not show a break may indicate that star formation is still occurring. Chandra observations have measured the breaks in the LFs of several nearby galaxies, e.g. M81 (Tennant et al. 2001), NGC 1553 (Blanton et al. 2001), NGC 4697 (Sarazin et al. 2001) and M83 (Soria & Wu 2002), and these breaks have been interpreted as evidence for impulsive star formation. Within M31, we find that the LF of three regions we studied have breaks at a different luminosities. The inner bulge has this break at the lowest luminosity, and the luminosity of the break increases monotonically as we go out from the inner bulge. If the breaks do indicate the epochs of star formation events, then these events occurred most recently in the disk of M31 and further back in time as we move towards the nucleus of M31.

As well as the monotonic shift in the break luminosity, there is a monotonic shift in the slopes of the LFs. As we move in towards the nucleus these slopes become progressively flatter. This is somewhat difficult to understand in the context of the discussion above, because the most luminous sources would be expected to have the shortest lifetimes. Loss of these sources as they age would tend to steepen the luminosity function, but we find flatter luminosity functions in the apparently older populations.

It is interesting to compare the LF of these three regions of M31 to those of other galaxies. M31 is not the first galaxy found to show a break in its LF nor is it the first to show different LFs in different regions; both M81 (Tennant et al. 2001) and M83 (Soria & Wu 2002) show similar behavior. In cases where a single slope is fit to the LF (ie, there is no clear break) the LFs of early type galaxies and the bulges of spirals tend to be steeper ($\alpha \sim 1.7$) than those of spiral disks and galaxies with active star forming regions ($\alpha \sim 0.8$, e.g, Prestwich 2001). The opposite seems to be the case within M31: the disk has a steeper LF than the bulge region. We speculate that this difference may be related to the location of the breaks in the M31 LF, which are at a lower luminosity that those seen in other galaxies. At these lower luminosities we may be sampling a different class of source, and the steepness of the LF may be due inclusion of this new class of faint sources rather than a loss of bright sources. We note that there is some evidence that we are sampling a different class of sources as we move out from the bulge, in that the fraction of sources which show variability decreases monotonically. If these sources have an intrinsically steeper LF than bright accreting binaries they may be able to cause the steepening of the LFs as we move from the bulge towards the disk.

7. CONCLUSIONS

In the past two years, *Chandra* and *XMM-Newton* have produced fruitful results on M31 (there are 7 papers related to M31 in this meeting). Clearly, it is just a beginning while data are still being analyzed and will be coming in in near future. A series of papers about the point source properties, the diffuse emission, the X-ray emission from the central supermassive black hole, and supersoft source populations are being prepared by our *Chandra* team. On the other hand, an extensive *XMM-Newton* survey on M31 is undergoing and the results would allow us for a detailed studies of point source properties and populations on the whole galaxy.

ACKNOWLEDGEMENTS

We are grateful to Kinwah Wu and Andrea Prestwich for stimulating discussion and comments. We acknowledge the support of a Croucher Fellowship (AKHK), NASA LTSA Grant NAG5-10889 (MRG,AKHK) and NAG5-10705 (RD,AKHK), and NASA Contract NAS8-39073 (MRG). The HRC GTO program is supported by NASA Contract NAS-38248.

REFERENCES

Berghoefer, T.W., Schmitt, J.H.M.M., Danner, R., Cassinelli, J.P. 1997, A&A, 322, 167

Blair, W.P., Kirshner, R.P., Chevalier, R.A. 1981, ApJ, 247, 879

Blanton, E.L., Sarazin, C.L., Irwin, J.A. 2001, ApJ, 552, 106

Chen, W, Shrader, C.R., Livio, M. 1997, ApJ, 491, 312

Crawford, D.F., Jauncey, D.L., Murdoch, H.S. 1970, ApJ, 162, 405

Di Stefano, R., Kong, A.K.H., Garcia, M.R., Barmby, P., Greiner, J., Murray, S.S., Primini, F.A. 2002, ApJ, in press (astro-ph/0106254)

d'Odorico, S., Dopita, M.A., Benvenuti, P. 1980, A&AS, 40, 67

Freeman, P.E., Kashyap, V., Rosner, R., Lamb, D.Q. 2002 ApJS, 138, 185

Garcia, M.R., Murray, S.S., Primini, F.A., Forman, W.R., McClintock, J.E., Jones, C. 2000a, ApJ, 537, L23

Garcia, M.R., Murray, S.S., Primini, F.A., McClintock, J.E., Callanan, P.J. 2000b, IAUC 7498

Garcia, M.R., Murray, S.S., Primini, F.A., Forman, W.R., Jones, C., McClintock, J.E. 2001a, in proceeding of IAU205, Galaxies at the Highest Angular Resolution (astro-ph/0012387)

Garcia, M., Kong, A., Primini, F., McClintock, J., Murray, S., Di stefano, R. 2001b, ATel 79

Garcia, M.R. 2001, Talk at 2 Years of Science with Chandra

Haas, M., Lemke, D., Stickel, M., Hippelein, H., Kunkel, M., Herbstmeier, U., Mattila, K. 1998, A&A, 338, L33

Hasinger, G., van der Klis, M. 1989, A&A, 225, 79

Kastner, J.H., Vrtilek, S.D., Soker, N. 2001, ApJ, 550, L189

Kong, A., Garcia, M., Murray, S., Primini, F., McClintock, J., Di Stefano, R. 2001, ATel 76

Kong, A.K.H., Charles, P.A. 2000, in Stellar Astrophysics, Proceedings of the Pacific Rim Conference: Kluwer Academic Publishers, p. 297

Osborne, J.P., et al. 2001, A&A, 378, 800

Prestwich, A.H. 2001, in Proceedings "The High Energy Universe in Sharp Focus", in press, ASP, eds. S. Vrtilek, E.M. Schegel & L. Kuhi (astro-ph/0108523)

Primini, F.A., Forman, W., Jones, C. 1993, ApJ, 410, 615

Primini, F., Garcia, M., Murray, S., Forman, W., Jones, C., McClintock, J. 2000, in proceedings 232 WE-Heraeus Seminar, 22-25 May 2000, Bad Honnef, Germany. Edited by Elly M. Berkhuijsen, Rainer Beck, and Rene A. M. Walterbos. Shaker, Aachen, p. 145-148

Psaltis, D., Lamb, F.K., Miller, G. S. 1995, ApJ, 454, L137

Sarazin, C.L., Irwin, J.A., Bregman, J.N. 2001, ApJ, 556, 533

Shirey, R., et al. 2001, A&A, 365, L195

Shirey, R. 2001, IAUC 7659

Smale, A.P., Kuulkers, E. 2000, ApJ, 528, 702

Soria, R., Wu, K. 2002, A&A, in press (astro-ph/0201059)

Supper, R., Hasinger, G., Lewin, W.H.G., Magnier, E.A., van Paradijs, J., Pietsch, W., Read, A.M., Trumper, J. 2001. A&A, 373, 63

Tennant, A.F., Wu, K., Ghosh, K.K., Kolodziejczak, J.J., Swartz, D.A. 2001, ApJ, 549, L43

Trudolyubov, S., Priedhorsky, W., Borozdin, K., Mason, K., Cordova, F. 2002 IAUC 7798

Wu, K., Tennant, A.F., Swartz, D.A., Ghosh, K.K., Hunstead, R.W. 2002, ApJ, submitted